

## Checklist of the Albanian mollusc fauna

by ZOLTÁN FEHÉR and ZOLTÁN PÉTER ERÖSS,  
Budapest

[2nd version, updated 28 August, 2009]

Taxa	Indicated by Fauna Europaea		Indicated by DHORA & WELTER- SCHULTES for Albania
	for Albania	for any European countries	
Fauna Europaea nomenclature is used generally, except for some special cases, when FE names are indicated in square brackets. * Indicates doubtful names and doubtful or incorrect Albanian occurrences			
<b>Unionidae</b>			
<i>Anodonta anatina</i> (LINNAEUS, 1758)	—	+	+
<i>Anodonta cygnea</i> (LINNAEUS, 1758)	+	+	+
* <i>Anodonta cymbalica</i> DROUET, 1879	—	—	* <sup>1</sup>
<i>Microcondylaea bonelli</i> (FÉRUSSAC, 1827) <sup>2</sup> [ <i>M. compressa</i> (MENKE, 1828)]	+	+	+
<i>Unio crassus carneus</i> KÜSTER, 1854	+	+	+
<i>Unio elongatulus pallens</i> ROSSMÄSSLER, 1842	+	+	+
<i>Unio elongatulus mancus</i> LAMARCK, 1819	—	+	+ <sup>3</sup>
<i>Unio pictorum</i> (LINNAEUS, 1758)	—	+	+
<i>Unio tumidus</i> PHILIPPSON, 1788 [ <i>U. tumidus</i> RETZIUS, 1788] <sup>4</sup>	—	+	* <sup>5</sup>
<b>Sphaeriidae</b>			
<i>Musculium lacustre</i> (O. F. MÜLLER, 1774)	—	+	+
* <i>Sphaerium parenzani</i> (GAMBETTA, 1930)	—	—	* <sup>6</sup>
<i>Pisidium casertanum</i> (POLI, 1791)	—	+	+
<i>Pisidium moitessierianum</i> PALADILHE, 1866 <sup>7</sup>	—	+	—
<i>Pisidium personatum</i> MALM, 1855 <sup>8</sup>	—	+	—
<i>Pisidium subtruncatum</i> MALM, 1855	—	+	+
<b>Dreissenidae</b>			
<i>Dreissena blanca</i> WESTERLUND, 1890 <sup>9</sup>	—	—	+
<i>Dreissena presbensis</i> KOBELT, 1915 <sup>10</sup>	—	—	+ <sup>11</sup>

<sup>1</sup>"systematic position not known"

<sup>2</sup> See FALKNER et al. (2001: 66).

<sup>3</sup> Mentioned as *U. mancus*.

<sup>4</sup> See FALKNER et al. (2001: 67).

<sup>5</sup> "Lives in Shkodra lake" - it means probably, that DHORA & WELTER-SCHULTES (1996) presumed its Albanian occurrence, but they were not aware of any verified Albanian record

<sup>6</sup> "systematic position not clear"

<sup>7</sup> First Albanian record: REISCHÜTZ et al. (2008b)

<sup>8</sup> First Albanian record: FEHÉR & ERÖSS (2009)

<sup>9</sup> This species is mentioned from the Shkodra Lake by DHORA & WELTER-SCHULTES (1996). ALBRECHT et al. (2007) have proved the distinctness of this species by molecular phylogenetic method, however they think that *D. blanca* lives not in the Shkodra Lake but in Prespa, Pamvotis, Trichonis and Amvrakia Lakes.

<sup>10</sup> According to ALBRECHT et al. (2007), *D. stankovici* LVOVA & STAROBOGATOV, 1982 is a synonym for this. Its distinctness has been proven by molecular phylogenetic method (e.g. GELEMBIUK et al. 2006). ALBRECHT et al. (2007) have found that this species occurs in the Shkodra, Ohrid, Dojran, Vegoritis, Prespa and Pamvotis Lakes (in the latter two syntopically with *D. blanca*).

<sup>11</sup> The Ohrid population is mentioned as *D. polymorpha* (PALLAS, 1771) by DHORA & WELTER-SCHULTES (1996).

Neritidae			
Theodoxus fluviatilis (Linnaeus, 1758)	+	+	+
Theodoxus varius (Menke, 1828)	+	+	+
Cochlostomatidae			
Cochlostoma scalarinum (A. & J.B. Villa, 1841) <sup>12</sup>	+	+	* <sup>13</sup>
Cochlostoma gracile gracile (L. Pfeiffer, 1849) <sup>14</sup>	—	+	
Cochlostoma gracile martensianum (Möllendorff, 1873)	+	+	+
Cochlostoma kleciaki arnautororum (A.J. Wagner, 1906)	+	+	+
Cochlostoma sturanii mnelense (A.J. Wagner, 1914)	+	+	+ <sup>15</sup>
Cochlostoma sturanii scalariniformis (A.J. Wagner, 1906)	+	+	
*Cochlostoma tergestinum (Westerlund, 1878) <sup>16</sup>	+	+	—
Cochlostoma auritum lamellatum (Westerlund, 1885) [C. a. alatum (A.J. Wagner, 1906)] <sup>17</sup>	+	+	+
Cochlostoma auritum gjonii Fehér, 2004	—	—	—
Cochlostoma georgi (A.J. Wagner, 1906)	+	+ <sup>18</sup>	+
Cochlostoma hoyeri hoyeri (Polinski, 1922)	+	+	+
Cochlostoma hoyeri csikii Fehér, 2004	—	—	—
Cochlostoma hoyeri lillae Fehér, 2004	—	—	—
Cochlostoma fuchsi Fehér, 2004	—	—	—
Cochlostoma pinteri pinteri Fehér, 2004	—	—	—
Cochlostoma pinteri hanswagneri Fehér, 2004	—	—	—
Cochlostoma pinteri erossi Fehér, 2004	—	—	—
Cochlostoma roseoli roseoli (A.J. Wagner, 1901)	+	+	
Cochlostoma roseoli kiriense (A.J. Wagner, 1906)	+	+	
Cochlostoma roseoli prekalense (Polinski, 1924)	+	+	+
Cochlostoma roseoli rioliense (A.J. Wagner, 1914)	+	+	
Cochlostoma roseoli scutariense (A.J. Wagner, 1906)	+	+	
Cochlostoma tessellatum tessellatum (Rossmässler, 1837) <sup>19</sup>	+	+	
Cochlostoma tessellatum excisum (Mousson, 1859)	+	+	+
Cochlostoma tessellatum tepelenum Fehér, Erőss & Varga, 2001	+	+	—
Aciculidae			
Platyla banatica (Rossmässler, 1842)	+	+	+
Platyla corpulenta Subai, 2009	—	—	—
Platyla pinteri (Subai, 1976)	+	+	+
Platyla wilhelmi (A.J. Wagner, 1910)	+	+	+
Renea kobelti albanica Boeters, Gittenberger & Subai, 1989	+	+	+
Viviparidae			
Viviparus contectus (Millet, 1813)	+	+	+
*Viviparus janinensis (Mousson, 1859)	—	+ <sup>20</sup>	* <sup>21</sup>
Viviparus mammillatus (Küster, 1852)	+	+	+
Viviparus viviparus (Linnaeus, 1758)	+	+	+

<sup>12</sup> First Albanian record: Fehér et al. (2001)

<sup>13</sup> "Lives probably not in today's Albania"

<sup>14</sup> Fehér et al. (2001) reported on Albanian populations that look more like the nominate subspecies.

<sup>15</sup> Mentioned as *C. sturanyi*.

<sup>16</sup> Most probably does not live in Albania.

<sup>17</sup> For nomenclatural notes see Fehér (2004).

<sup>18</sup> Despite Fauna Europaea lists this species also from Yugoslavia, it is known only from Albania.

<sup>19</sup> First Albanian record: Fehér et al. (2001).

<sup>20</sup> Synonym for *V. mammillatus*.

<sup>21</sup> "occurrence in Albania needs confirmation"

Melanopsidae			
Amphimelania holandrii (C. Pfeiffer, 1828)	+	+	+
Pomatidae			
Pomatias elegans (O.F. Müller, 1774)	+	+	+
Bithyniidae			
*Bithynia tentaculata (Linnaeus, 1758) <sup>22</sup>	+	+	+
Bithynia radomani Glöer & Pesić, 2007 <sup>23</sup>	—	—	—
*Bithynia montenegrina (Wohlberedt, 1901) <sup>24</sup>	—	—	—
*Bithynia schwabii Frauenfeld, 1865 <sup>25</sup>	+	+	—
*Bithynia walderdorffii Frauenfeld, 1865 <sup>26</sup>	+	+	—
Hydrobiidae			
Adriohydrobia gagatinella (Küster, 1852)	+	+	* <sup>27</sup>
Belgrandia ionica (Schütt 1980) <sup>28</sup>	—	+	—
Bythinella cf. drimica Radoman, 1976 <sup>29</sup>	—	+	—
Grossuana euxina (A.J. Wagner, 1828) <sup>30</sup>	—	+	—
Horatia klecakiana Bourguignat, 1887	—	+	* <sup>31</sup>
*Hydrobia ventrosa (Montagu, 1803)	—	+	+ <sup>32</sup>
*Hydrobia vitrea (Risso, 1826)	+	+	—
Malaprespa albanica Radoman, 1973	+	+	+ <sup>33</sup>
Ohridohoratia polinskii (Radoman, 1960)	+	+	+ <sup>34</sup>
Ohridohoratia pygmaea (Westerlund, 1902)	+	+	+
Ohridohoratia sturanyi (Westerlund, 1902) <sup>35</sup>	—	+	—
Ohrigocea stankovici (Hadžišće, 1959)	+	+	+
Radomaniola curta (Küster, 1853) <sup>36</sup>	—	+	+ <sup>37</sup>
Radomaniola albanica Radoman, 1973 <sup>38</sup>	+	+	
Paladilhiopsis cf. serbica (Pavlovic, 1913) <sup>39</sup>	—	+	—
Parabythinella malaprespensis Radoman, 1973	+	+	+

<sup>22</sup> Probably all records refer to *B. radomani*.

<sup>23</sup> First Albanian record: Fehér & Erőss (2009)

<sup>24</sup> Mentioned by Reischütz et al. (2008b), probably refers to *B. radomani*.

<sup>25</sup> Probably does not live in Albania.

<sup>26</sup> Probably does not live in Albania.

<sup>27</sup> Mentioned as *A. consociella* with the note "occurrence in Albania needs confirmation". There is a lot in the collection of the Hungarian Natural History Museum, collected in Himara (South Albania) by the Albanian entomologist X. Murraj, which is labelled as "*Adriohydrobia consociella*", but it was revised by M. Bodon and proved to be *Radomaniola albanica*.

<sup>28</sup> First Albanian record: Fehér & Erőss (2009).

<sup>29</sup> First Albanian record: Fehér & Erőss (2009).

<sup>30</sup> = *G. serbica* sensu Radoman. First Albanian record: Fehér & Erőss (2009).

<sup>31</sup> "Lives probably not in today's Albania"

<sup>32</sup> Dhora & Welter-Schultes (1996) reported this species from three Albanian localities. Two of them are inland and therefore, these records are surely incorrect, since *H. ventrosa* lives always in brackish water. The third locality is mapped in the coast, thus this record is not impossible, but needs confirmation (M. Bodon personal communication).

<sup>33</sup> Dhora & Welter-Schultes (1996) indicate *Albaniana albanica* as a separate species, according to the Fauna Europaea, *M. albanica* and *A. albanica* are synonyms.

<sup>34</sup> Mentioned as Polinskiola polinskii

<sup>35</sup> First Albanian record: Reischütz et al. (2008b), see main text for further notes and distribution data.

<sup>36</sup> First Albanian record: Fehér & Erőss (2009), see main text for details.

<sup>37</sup> Mentioned by Dhora & Welter-Schultes (1996) as *Orientalina*, but being a junior homonym, that name must be replaced (see main text for details). Dhora & Welter-Schultes (1996) treat "*Orientalina albanica*" as the subspecies of "*O. curta*" and they map only the published "*O. albanica*" records.

<sup>38</sup> Possibly a synonym for "*Hydrobia haesitans* Westerlund, 1881", see the main text for details.

<sup>39</sup> First Albanian record: Fehér & Erőss (2009). See the text for details.

Prespolitoria malaprespensis Radoman, 1973	+	+	+
Pseudohoratia ohridana (Polinski, 1929) <sup>40</sup>	—	+	—
Vinodolia (Anagastina) scutarica (Radoman, 1973)	+	+	+
Zaumia kusceri (Hadžišče, 1959)	+	+	+
Emmericiidae			
*Emmericia patula (Brumati, 1838)	—	+	* <sup>41</sup>
Pyrgulidae			
Ochridopyrgula macedonica (Brusina, 1896)	+	+	+
Chilopyrgula sturanyi (Brusina, 1896)	+	+	+
Pyrgula annulata dalmatica Schütt, 1968	+	+	+ <sup>42</sup>
Pyrgohydrobia grochmalickii (Polinski 1929) <sup>43</sup>	—	+	—
Valvatidae			
Valvata cristata O.F. Müller, 1774	+	+	+
Valvata piscinalis (O.F. Müller, 1774)	+	+	+
Valvata rhabdota Sturany, 1894	—	+	* <sup>44</sup>
Valvata relictia (Polinski, 1929) <sup>45</sup>	—	+	—
Valvata stenotrema Polinski, 1929 <sup>46</sup>	—	+	—
Carychiidae <sup>47</sup>			
Carychium tridentatum (Risso, 1826) <sup>48</sup>	—	+	—
Carychium minimum O.F. Müller, 1774 <sup>49</sup>	—	+	—
Ellobiidae			
Myosotella myosotis (Draparnaud, 1801)	—	+	* <sup>50</sup>
Acroloxidae			
Acroloxus lacustris (Linnaeus, 1758)	+	+	+
Lymnaeidae			
Lymnaea stagnalis (Linnaeus, 1758)	+	+	+
Radix auricularia (Linnaeus, 1758)	+	+	+
Radix ampla (W. Hartmann, 1821)	+	+	+
Radix balthica (Linnaeus, 1758)	+	+	—
Radix labiata (Rossmässler, 1835)	+	+	+ <sup>51</sup>
Radix relictia Polinski, 1929	+	+	—
Stagnicola corvus (Gmelin, 1791)	+	+	+
Stagnicola palustris (O.F. Müller, 1774)	+	+	+
Galba truncatula (O.F. Müller, 1774)	+	+	+
Physidae			
Aplexa hypnorum (Linnaeus, 1758)	+	+	+
Physella acuta (Draparnaud, 1805)	+	+	+

<sup>40</sup> First Albanian record: Fehér & Drimmer (2004).

<sup>41</sup> "occurrence in Albania needs confirmation"

<sup>42</sup> Mentioned as *P. annulata*.

<sup>43</sup> First Albanian record: Fehér & Erőss (2009).

<sup>44</sup> "occurrence in Albania needs confirmation"

<sup>45</sup> First Albanian record: Fehér & Erőss (2009).

<sup>46</sup> First Albanian record: Fehér & Erőss (2009).

<sup>47</sup> *Carychium* spp. have not been known from Albania to now, first records are presented in this study. Our records are from the northern part of Albania. Since *C. hellenicum* Bank & Gittenberger, 1985 is known from Epirus and Korfu Island, this species probably lives in South Albania.

<sup>48</sup> First Albanian record: Fehér & Erőss (2009).

<sup>49</sup> Reischütz et al. (2008b) have found subrecent shells. First verified Albanian record: Fehér & Erőss (2009).

<sup>50</sup> "occurrence in Albania needs confirmation". See the main text for a recent distribution record.

<sup>51</sup> Mentioned as *R. peregra*.

<i>Physa fontinalis</i> (Linnaeus, 1758)	+	+	+
<b>Planorbidae</b>			
<i>Ancylus fluviatilis</i> O.F. Müller, 1774	+	+	+
<i>Ancylus tapirulus</i> Polinski, 1929 <sup>52</sup>	—	+	—
<i>Gyraulus acronicus</i> (A. Féruccac, 1807)	+	+	—
<i>Gyraulus albidus</i> Radoman, 1953 <sup>53</sup>	—	+	—
<i>Gyraulus albus</i> (O.F. Müller, 1774)	+	+	+
<i>Gyraulus laevis</i> (Alder, 1838)	+	+	+
<i>Gyraulus crista</i> (Linnaeus, 1758)	+	+	+
<i>Gyraulus crenophilus</i> Hubendick & Radoman, 1959	+	+	+
<i>Gyraulus fontinalis</i> Hubendick & Radoman, 1959	+	+	+
<i>Gyraulus trapezoides</i> Polinski, 1929 <sup>54</sup>	—	+	—
<i>Bathyomphalus contortus</i> (Linnaeus, 1758)	+	+	+
* <i>Anisus vortex</i> (Linnaeus, 1758)	+	+	+ <sup>55</sup>
* <i>Anisus vorticulus</i> (Troschel, 1834)	+	+	+ <sup>56</sup>
<i>Anisus spirorbis</i> (Linnaeus, 1758) <sup>57</sup>	—	+	—
<i>Planorbis carinatus</i> O.F. Müller, 1774	+	+	+
<i>Planorbis planorbis</i> (Linnaeus, 1758)	+	+	+
<i>Planorbis presbensis</i> Sturany, 1894 <sup>58</sup>	—	+	—
<i>Ferrissia clessiniana</i> (Jickeli, 1882)	+	+	+ <sup>59</sup>
<i>Planorbarius corneus</i> corneus (Linnaeus, 1758)	+	+	+
<i>Planorbarius corneus</i> cf. <i>grandis</i> (Dunker, 1850) <sup>60</sup>	—	+	
<i>Planorbarius corneus</i> <i>arabatzis</i> Reischütz, Reischütz & Fischer, 2008 <sup>61</sup>	—	—	
* <i>Planorbella anceps</i> (Menke, 1830) <sup>62</sup>	—	—	—
<b>Succineidae</b>			

<sup>52</sup> Fauna Europaea list this species only from Macedonia. First Albanian record: Reischütz et al. (2008b). Albrecht et al. (2006) studied recently the molecular phylogeny of the *Ancylus* species of the Lake Ohrid, mostly based on Macedonian material. They have found that *A. fluviatilis* does not occur in the lake, just *A. tapirulus*, *A. scalariformis*, *A. lapicidus* and an undescribed fourth *Ancylus* species.

<sup>53</sup> First Albanian record: Fehér & Erőss (2009).

<sup>54</sup> First Albanian record: Fehér & Erőss (2009).

<sup>55</sup> Listed by Dhora & Welter-Schlüter (1996) without any distribution map and specific distribution data. According to their reference list, Dhora & Welter-Schlüter (1996) included this species in the Albanian fauna list on the basis of a roughly drawn distribution map of Meier-Brook (1983: p. 92).

<sup>56</sup> Listed by Dhora & Welter-Schlüter (1996) without any distribution map and specific distribution data. According to their reference list, Dhora & Welter-Schlüter (1996) included this species in the Albanian fauna list on the basis of a roughly drawn distribution map of Meier-Brook (1983: p. 92).

<sup>57</sup> First Albanian record: Reischütz et al. (2008b).

<sup>58</sup> First Albanian records: Fehér & Drimmer (2004) and Fehér & Erőss (2009).

<sup>59</sup> Mentioned as *F. wautieri*.

<sup>60</sup> According to the Fauna Europaea, *P. c. grandis* occurs only in Greece. In the Prespa Lakes, *P. corneus* populations are significantly different from the nominate form and resemble to the figure in the original description of *Planorbis grandis*. Reischütz et al. (2008a) believes that two distinct subspecies occur in the Prespa region: *P. corneus grandis* in the Small Prespa Lake and *P. corneus arabatzis* in the Great Prepa Lake. The distinction of their new subspecies was based on slight shell morphological differences. As the two lakes are divided only by a narrow isthmus and they are connected by a creek, it can be supposed reasonably that *P. corneus* populations of the two lakes are in connection. In view of the above, it is questionable and needs further confirmation whether or not the the Great Prespa populations belong to a separate taxon.

<sup>61</sup> See the above note.

<sup>62</sup> The abnormal *P. corneus* form in the Great Prespa Lake, was misidentified and incorrectly reported as *P. anceps* by Erőss et al. (2005). See the main text for details.

Oxyloma (Oxyloma) elegans (Risso, 1826)	+	+	+
*Succinella oblonga (Draparnaud, 1801)	+	+	+ (!) <sup>64</sup>
Cochlicopidae			
Cochlicopa lubrica (O.F. Müller, 1774)	+	+	+
Cochlicopa lubricicella (Porro, 1838)	+	+	+
Azecidae			
Hypnophila pupaeformis (Cantraine, 1835)	+	+	+
Hypnophila polita (Porro, 1838)	—	+	* <sup>65</sup>
Lauriidae			
Lauria cylindracea (Da Costa, 1778)	+	+	+
Lauria sempronii (Charpentier, 1837) <sup>66</sup>	—	+	—
Orculidae			
Pagodulina kaeufeli Klemm, 1939	+	+	+
Pagodulina subdola gracilior Pilsbry, 1926	+	+	+
Sphyradium doliolum (Bruguière, 1792)	+	+	+
Orcula schmidti schmidti (Küster, 1843)	+	+	+
Orcula wagneri wagneri Sturany, 1914	+	+	+
Argnidae			
Agardhiella truncatella (L. Pfeiffer, 1841)	—	+	
Agardhiella skipetarica <sup>69</sup> (A.J. Wagner, 1914) [A. truncatella skipetarica]	—	+	+
*Agardhiella lamellata (Clessin, 1887) <sup>70</sup>	+	+	—
Agardhiella zoltanorum Subai, 2008	—	—	—
Agardhiella extravaganta Subai, 2008	—	—	—
Strobilopsidae			
Spelaeodiscus albanicus (A.J. Wagner, 1914)	+	+	+
Valloniidae			
Vallonia costata (O.F. Müller, 1774)	+	+	+
Vallonia enniensis (Gredler, 1856)	+	+	+
Vallonia excentrica Sterki, 1893	+	+	+
Vallonia pulchella (O.F. Müller, 1774)	+	+	+
Acanthinula aculeata (O. F. Müller, 1774) <sup>71</sup>	—	+	* <sup>72</sup>
Gittenbergia sororcula (Benoit, 1859) <sup>73</sup>	—	+	* <sup>74</sup>
*Plagyrone placida (Shuttleworth, 1852) <sup>75</sup>	—	+	—
Pupillidae			

<sup>63</sup> Dhora & Welter-Schultes (1996) list *Succinea pfeifferi* Rossmassler, 1835 separately, which is however a synonym for this species.

<sup>64</sup> Due to a misinterpretation of Bába (1982), this species was falsely included in the Albanian list by Dhora & Welter-Schultes (1996).

<sup>65</sup> "lives very probably in Albania"

<sup>66</sup> First Albanian record: Fehér & Erőss (2009).

<sup>67</sup> Mentioned as *P. subdola kaeufeli*.

<sup>68</sup> Dhora & Welter-Schultes (1996) indicates this two species as "syn/ssp.". In their map (p. 120) only one locality (the type locality of *A. skipetarica*) is indicated.

<sup>69</sup> According to Subai (1999a and 2008) this is a distinct species.

<sup>70</sup> Does not live in Albania. Found in Romania and in Hungary (fluvial flotsam) up to now.

<sup>71</sup> First Albanian record: Fehér & Erőss (2009).

<sup>72</sup> "occurrence in Albania needs confirmation"

<sup>73</sup> First Albanian record: Fehér & Erőss (2009).

<sup>74</sup> "occurrence in Albania needs confirmation"

<sup>75</sup> Mentioned by Reischütz et al. (2008b).

* <i>Pupilla alpicola</i> (Charpentier, 1837) <sup>76</sup>	+	+	+
* <i>Pupilla triplicata</i> (S. Studer, 1820)	+	+	+ (!) <sup>77</sup>
<i>Pupilla sterrii</i> (Voith, 1840) <sup>78</sup>	—	+	—
Pyramidulidae			
<i>Pyramidula cephalonica</i> (Westerlund, 1898)	+	+	+
<i>Pyramidula pusilla</i> (Vallot, 1801)	+	+	+
Chondrinidae			
<i>Granopupa granum</i> (Draparnaud, 1801)	+	+	+
* <i>Granaria frumentum illyrica</i> (Rossmässler, 1835) [G. illyrica]	+ <sup>79</sup>	+	
<i>Granaria frumentum hungarica</i> (M. von Kimakowicz, 1890)	—	+ <sup>80</sup>	+
<i>Granaria frumentum</i> ssp. n. <sup>81</sup>	—	—	
<i>Chondrina arcadica</i> abundans (Westerlund, 1894)	+	+	
<i>Chondrina arcadica</i> clienta (Westerlund, 1883)	+	+	+
<i>Chondrina spelta</i> spelta (Beck, 1837) <sup>82</sup>	—	+	
<i>Chondrina spelta</i> fuchsii H. Nordsieck, 1970	+	+	
<i>Chondrina spelta</i> serbica H. Nordsieck, 1970	+	+	
<i>Rupestrella philippii</i> (Cantraine, 1841)	+	+	+
<i>Rupestrella rhodia</i> (Roth, 1839)	+	+	+
Vertiginidae			
<i>Vertigo pusilla</i> O.F. Müller, 1774 <sup>83</sup>	+	+	+ (!) <sup>84</sup>
<i>Vertigo pygmaea</i> (Draparnaud, 1801) <sup>85</sup>	—	+	—
* <i>Vertigo antivertigo</i> (Draparnaud, 1801) <sup>86</sup>	—	+	—
<i>Columella edentula</i> (Draparnaud, 1805) <sup>87</sup>	—	+	—
<i>Truncatellina callicratidis</i> (Scacchi, 1833)	+	+	+
<i>Truncatellina claustralis</i> (Gredler, 1856)	+	+	+
<i>Truncatellina cylindrica</i> (A. Féruccac, 1807)	+	+	+
<i>Truncatellina rothi</i> (Reinhardt, 1916)	+	+	+
Enidae			
<i>Ena concolor</i> (Westerlund, 1887)	+	+	+
<i>Ena montana</i> (Draparnaud, 1801)	+	+	+
<i>Ena subtilis subtilis</i> (Rossmässler, 1837)	+	+	
<i>Ena subtilis coryrensis</i> (Mousson, 1859)	+	+	+
<i>Jaminia quadridens</i> (O.F. Müller, 1774)	+	+	+

<sup>76</sup> Soós (1924) reports on the occurrence of this species in the Korab Mts., that record, however, possibly based on misidentified *P. sterrii* material.

<sup>77</sup> Due to a misinterpretation of Bába (1982), this species was included in the Albanian list by Dhora & Welter-Schultes (1996), but there are no confirmed Albanian record up to now.

<sup>78</sup> First Albanian record: Fehér & Erőss (2009).

<sup>79</sup> Fauna Europaea treats this taxon as a distinct species.

<sup>80</sup> Fauna Europaea treats this taxon as a synonym for the nominate subspecies.

<sup>81</sup> According to a recent revision of the Balkan species of *Granaria*, "G. illyrica" is the subspecies of *G. frumentum*, *G. f. hungarica* is a valid subspecies and only two *G. frumentum* subspecies occur in Albania: *G. f. hungarica* and a new form (Fehér et al. submitted).

<sup>82</sup> Although Fauna Europaea does not indicate it, the nominate subspecies lives in Albania too (see Nordsieck 1970).

<sup>83</sup> First verified Albanian records are given in the Fehér & Erőss (2009).

<sup>84</sup> This species was included in the Albanian checklist of Dhora & Welter-Schultes (1996) due to a misinterpretation of Bába (1982).

<sup>85</sup> First Albanian record: Reischütz et al. (2008b), further distribution data are given in the Fehér & Erőss (2009).

<sup>86</sup> A subrecent occurrence was reported by Reischütz et al. (2008b), recent occurrence needs confirmation.

<sup>87</sup> First Albanian record: Fehér & Erőss (2009).

Pseudochondrula seductilis (Rossmässler, 1837)	+	+	+
Multidentula cf. squalina (L. Pfeiffer, 1848) <sup>88</sup>	—	—	—
Merdigera obscura (O.F. Müller, 1774)	+	+	+
Mastus grandis grandis (Mousson, 1859)	+	+	+ <sup>89</sup>
Chondrula consentanea (Westerlund, 1887)	—	+	* <sup>90</sup>
Chondrula lugorensis A.J. Wagner, 1914	+	+	+
Chondrula macedonica A.J. Wagner, 1914	+	+	+
Chondrula microtragus (Rossmässler, 1839)	+	+	+
Chondrula quinquedentata (Rossmässler, 1837)	+	+	+ <sup>91</sup>
Chondrula tridens (O.F. Müller, 1774)	+	+	
Napaeopsis cephalonicus (Mousson, 1859)	+	—	+ <sup>92</sup>
Napaeopsis merditanus merditanus (Sturany, 1907)	+	+	
Napaeopsis merditanus latifianus (Sturany, 1907)	+	+	
Napaeopsis merditanus winneguthi (Sturany, 1907)	+	+	
Napaeopsis merditanus zebianus (Sturany, 1907)	+	+	
Zebrina detrita (O.F. Müller, 1774)	+	+	+
Zebrina detrita sallake ssp. n. <sup>93</sup>	—	—	—
Clausiliidae			
Sciocochlea illogaraensis (Reischütz & Reischütz, 2009)	—	—	—
Bulgarica vetusta gracilior S.H.F. Jaeckel, 1954	+	+	+
Bulgarica vetusta nannodes (A.J. Wagner, 1912)	+	+	
Vestia roschitzii apragmosyne (A.J. Wagner, 1919)	+	+	+
Alinda biplicata biplicata (Montagu, 1803) <sup>94</sup> [Balea biplicata]	—	+	+ <sup>95</sup>
Alinda biplicata byshekensis Nordsieck, 2008	—	—	
Alinda biplicata distincta (Sturany, 1894) <sup>95</sup>	+	+	
*Alinda biplicata labiata (Westerlund, 1885) <sup>96</sup>	+	+	
*Alinda biplicata metriotes (A.J. Wagner, 1919) <sup>97</sup>	+	+	
Balea serbica serbica (Möllendorff, 1873)	+	+	+ <sup>98</sup>
Laciniaria plicata (Draparnaud, 1801)	+	+	+
Clausilia dubia vindobonensis A. Schmidt, 1856	+	+	* <sup>99</sup>
Macrogaster ventricosa (Draparnaud, 1801) <sup>100</sup>	+	+	—
Strigilodelima conspersa (L. Pfeiffer, 1848)	— <sup>101</sup>	+	+
Strigilodelima pentheri (A.J. Wagner, 1919)	+	+	+
* Albinaria maculosa (Deshayes, 1835)	+	+	
[Albinaria s. schuchii (Rossmässler, 1836)] <sup>102</sup>	+	+	+

<sup>88</sup> First Albanian record: Fehér & Erőss (2009).

<sup>89</sup> Mentioned as *M. pupa*.

<sup>90</sup> "occurrence in Albania needs confirmation"

<sup>91</sup> Dhora & Welter-Schultes (1996) treat *C. quinquedentata* as a "syn/ssp." of *C. tridens*.

<sup>92</sup> Dhora & Welter-Schultes (1996) treat *merditanus*, *latifianus*, *winneguthi* and *zebianus* as "syn/ssp." of the *N. cephalonicus*. Fauna Europaea indicates *N. cephalonicus* only from Greece.

<sup>93</sup> Described in the Fehér & Erőss (2009).

<sup>94</sup> Systematics and distribution of this species has been revised by Nordsieck (2008).

<sup>95</sup> Though Dhora & Welter-Schultes (1996) do not go into the particulars at the subspecies level generally, but for this subspecies they note that "*B. b. distincta* does not live in Albania". But see Nordsieck (2008).

<sup>96</sup> According to Nordsieck (2008), this is a synonym of the nominate subspecies ("oreinos form").

<sup>97</sup> According to Nordsieck (2008), this is a synonym of the nominate subspecies ("oreinos form").

<sup>98</sup> Dhora & Welter-Schultes (1996) treat this taxon as a "syn/ssp." of *B. fallax*.

<sup>99</sup> "needs confirmation"

<sup>100</sup> First Albanian record: Dhora & Welter-Schultes (1999a).

<sup>101</sup> Fauna Europaea lists this species only for Macedonia. It is very common in Albania.

<sup>102</sup> Nordsieck (1974) indicates *A. schuchii* from Albania (collected near Sarande by Edlauer). Despite our efforts, we could not confirm this occurrence to now. Nordsieck (1995) proposed to use the name *schuchii*

<i>Albinaria scopulosa epirotea</i> H. Nordsieck, 1974	+	+	+
<i>Albinaria scopulosa winneguthi</i> (A.J. Wagner, 1914)	+	+	
<i>Albinaria senilis inconstans</i> (Mousson, 1859) <sup>103</sup>	—	+	—
<i>Agathylla biloba biloba</i> (A.J. Wagner, 1914)	+	+	+
<i>Agathylla biloba dabovici</i> Brandt, 1961	+	+	
<i>Agathylla biloba martae</i> Erőss & Szekeres, 1999	+	+	—
<i>Agathylla neutra neutra</i> (Westerlund, 1898)	+	+	+ <sup>104</sup>
<i>Agathylla neutra merditana</i> (A.J. Wagner, 1914)	+	+	+
<i>Medora proxima remota</i> (Polinski, 1924)	+	+	+
<i>Triloba sandrii</i> (Küster, 1844)	+	+	+
<i>Triloba thaumasia thaumasia</i> (Sturany, 1907)	+	+	+
<i>Triloba thaumasia dardanorum</i> (A.J. Wagner, 1919)	+	+	
<i>Triloba thaumasia korabensis</i> (A.J. Wagner, 1919)	+	+	
* <i>Herilla bosniensis</i> (L. Pfeiffer, 1868)	—	+	+ <sup>105</sup>
<i>Herilla illyrica diabasis</i> (A.J. Wagner, 1919)	+	+	+
<i>Herilla illyrica oribates</i> (Sturany, 1909)	—	+	
<i>Herilla jabucica interrupta</i> Nordsieck, 1971 <sup>106</sup>	—	+	—
<i>Herilla ziegleri klemmi</i> Nordsieck, 1971 <sup>107</sup>	—	+	+
<i>Protoherilla baleiformis</i> (O. Boettger, 1909)	+	+	+
<i>Protoherilla pseudofallax</i> (Nordsieck, 1972)	—	+	+
<i>Montenegrina apfelbecki apfelbecki</i> (Sturany, 1907)	+	+	+
<i>Montenegrina apfelbecki caesia</i> Fehér & Szekeres, 2006 <sup>108</sup>	—	—	—
<i>Montenegrina apfelbecki csikii</i> Erőss & Szekeres, 2006 <sup>109</sup>	—	—	—
<i>Montenegrina apfelbecki okolensis</i> Szekeres, 2006 <sup>110</sup>	—	—	—
<i>Montenegrina apfelbecki wagneri</i> Szekeres, 2006 <sup>111</sup>	—	—	—
<i>Montenegrina dofleini occidentalis</i> Nordsieck, 1977 <sup>112</sup>	—	+	—
<i>Montenegrina dofleini plenostoma</i> Fehér & Szekeres, 2006	—	—	—
<i>Montenegrina drimmeri</i> Fehér & Szekeres, 2006	—	—	—
<i>Montenegrina ersekensis ersekensis</i> H. Nordsieck, 1996 <sup>113</sup>	+	+	+
<i>Montenegrina ersekensis nobilis</i> Erőss & Szekeres, 2006 <sup>114</sup>	—	—	—

instead of *maculosa* because several authors had misunderstood the name *maculosa* before, and he considered it as a nomen oblitum, but this view is not tenable (the name was not forgotten, only misinterpreted). Since Nordsieck did not put his proposal into a frame of nomenclatural rules, we have to use the name *maculosa* which has priority (Welter-Schultes personal communication).

<sup>103</sup> First Albanian record: Fehér & Erőss (2009).

<sup>104</sup> Mentioned by Dhora & Welter-Schultes (1996) as *A. merditana scraparana* Nordsieck 1996. See Nordsieck (2001) for details.

<sup>105</sup> Dhora & Welter-Schultes (1996) lists this species without a distribution map. Though there are 3 subspecies indicated for Albania in the table of Jaeckel et al. (1957), but this is probably a misprint. To our present knowledge, *H. bosniensis* does not occur in Albania.

<sup>106</sup> First Albanian record: Dhora & Welter-Schultes (1999a).

<sup>107</sup> First Albanian record: Nordsieck (1971).

<sup>108</sup> Nordsieck (2007) mentions as the subspecies of *M. janinensis*.

<sup>109</sup> Nordsieck (2007) mentions as the subspecies of *M. janinensis*.

<sup>110</sup> Nordsieck (2007) mentions as the subspecies of *M. janinensis*, later, proposed as the synonym of "*M. j. caesia*" (Nordsieck 2009).

<sup>111</sup> Probably, in the original description (Erőss et al. 2006) the type locality (Mali Thate) was incorrectly identified with Maja e Madhe in North Albania. The type locality is more probably identical with Mali i Thatës in Southeast Albania. In Nordsieck's (2009) opinion, this taxon is the synonym of *M. janinensis fagorum*.

<sup>112</sup> First Albanian record: Fehér & Erőss (2009).

<sup>113</sup> Nordsieck (2007) mentions as the subspecies of *M. irmengardis*, and later (Nordsieck 2009) as the subspecies of *M. skipetarica*. See note at *M. perstriata skipetarica*.

<sup>114</sup> Its subspecies rank was questioned by Nordsieck (2009).

Montenegrina fuchsi fuchsi R.A. Brandt, 1961	+	+	+
Montenegrina fuchsi klemmi R.A. Brandt, 1962	+	+	+
Montenegrina fuchsi muranyii Feher & Szekeres, 2006	—	—	—
Montenegrina helvola helvola (Küster, 1853)	+	+	+
Montenegrina helvola pageti R.A. Brandt, 1962	+	+	
Montenegrina helvola carinata Erőss & Szekeres, 1999	+	+	—
Montenegrina helvola magna Fehér & Szekeres, 2006	—	—	—
Montenegrina helvola ornata Erőss & Szekeres, 1999	+	+	—
Montenegrina irmengardis flava Erőss & Szekeres, 2006	—	—	—
Montenegrina irmengardis remota Feher & Szekeres, 2006	—	—	—
Montenegrina irmengardis rugosa Feher & Szekeres, 2006	—	—	—
Montenegrina irmengardis timeae Erőss & Szekeres, 2006	—	—	—
Montenegrina irmengardis welterschultesi Feher & Szekeres, 1999	+	+	+ <sup>115</sup>
Montenegrina janinensis coerulescens H. Nordsieck, 1996	+	+	+
Montenegrina janinensis fagorum H. Nordsieck, 1974	+	+	
Montenegrina janinensis tomorosi R.A. Brandt, 1961	+	+	
Montenegrina janinensis ampla Feher & Szekeres, 2006	—	—	—
Montenegrina janinensis soosi Erőss & Szekeres, 2006	—	—	—
Montenegrina laxa laxa (Küster, 1860)	+	+	+
Montenegrina laxa chiasma H. Nordsieck, 1972	+	+	
Montenegrina laxa iba H. Nordsieck, 1972	+	+	
Montenegrina laxa miraka H. Nordsieck, 1996	+	+	
Montenegrina laxa disjuncta Fehér & Szekeres, 2006	—	—	—
Montenegrina laxa errans Erőss & Szekeres, 2006	—	—	—
Montenegrina laxa kontschani Erőss & Szekeres, 2006	—	—	—
Montenegrina minuscula Erőss & Szekeres, 2006	—	—	—
Montenegrina perstriata callistoma Fehér & Szekeres, 2006	—	—	—
Montenegrina perstriata crassa Erőss & Szekeres, 1999 <sup>116</sup>	—	+	—
Montenegrina perstriata gracilis Erőss & Szekeres, 2006	—	—	—
Montenegrina perstriata nana Fehér & Szekeres, 2006	—	—	—
Montenegrina perstriata skipetarica (Soós, 1924) <sup>117</sup>	+	+	+
Montenegrina perstriata tenebrosa Nordsieck, 2009	—	—	—
Montenegrina sattmanni sattmanni Nordsieck, 1988 <sup>118</sup>	—	+	—
Montenegrina sattmanni costulata Erőss & Szekeres, 2006	—	—	—
Montenegrina sattmanni fusca Fehér & Szekeres, 2006 <sup>119</sup>	—	—	—
Montenegrina subcristata wohlberedti (Möllendorff, 1899)	+	+	+
*Cochlodina commutata (Rossmässler, 1836) <sup>120</sup>	+	+	—
Cochlodina laminata albanica S.H.F. Jaeckel, 1956	+	+	+
Charpentieria lamellata (Rossmässler, 1836)	+	+	+ <sup>121</sup>
Charpentieria pantocratoris margaritifera (H. Nordsieck, 1996)	+	+	+

<sup>115</sup> The record of *M. irmengardis* by Dhora & Welter-Schultes (1996) was based on this subspecies.

<sup>116</sup> First Albanian record: Fehér & Erőss (2009). Nordsieck (2009) questionned its subspecies rank.

<sup>117</sup> In Nordsieck's (2009) opinion, this subspecies belongs to *M. irmengardis* rather than to *M. perstriata*. As *skipetarica* is an older name than *irmengardis*, he suggest to use this specific name for each *M. irmengardis* subspecies.

<sup>118</sup> First Albanian record: Erőss et al. (2006).

<sup>119</sup> In Nordsieck's (2009) opinion it belongs to "*M. skipetarica*" rather than to *M. sattmanni*. See note at *M. perstriata skipetarica*.

<sup>120</sup> Most probably does not live in Albania.

<sup>121</sup> Mentioned as *Siciliaria lamellata*.

Charpentieria splendens (H. Nordsieck, 1996)	+	+	+ <sup>122</sup>
Charpentieria stigmatica stigmatica (Rossmässler, 1836)	+	+	+
Charpentieria stigmatica sturmii (L. Pfeiffer, 1848)	+	+	+
Papillifera bidens (Linnaeus, 1758)	—	+	+ <sup>123</sup>
Delima bilabiata (J.A. Wagner, 1829)	—	+	+
Delima montenegrina (L. Pfeiffer, 1848)	—	+	+
*Delima binotata (Rossmässler, 1836)	+	+	* <sup>124</sup>
*Delima blanda (Rossmässler, 1836)	+	+	—
Ferussaciidae			
Cecilioides acicula (O.F. Müller, 1774)	+	+	+
Cecilioides tumulorum (Bourguignat, 1856)	+	+	+
Subulinidae			
*Rumina decollata (Linnaeus, 1758)	+	+	* <sup>125</sup>
Oleacinidae			
Poiretia compressa (Mousson, 1859)	+	+	+
Poiretia cornea (Brumati, 1838)	+	+	+
Poiretia delesserti (Bourguignat, 1852)	+	+	+
Helicodiscidae			
Lucilla singleyana (Pilsbry, 1889) <sup>126</sup>	—	+	—
Punctidae			
Punctum pygmaeum (Draparnaud, 1801)	+	+	+
Euconulidae			
Euconulus fulvus (O.F. Müller, 1774)	+	+	+
Gastodontidae			
Zonitoides nitidus (O.F. Müller, 1774)	+	+	+
Zonitidae [incl. Pristilomatidae, Oxychilidae, Zonitidae]			
Vitrella botterii (L. Pfeiffer, 1853)	+	+	+
Vitrella contracta (Westerlund, 1871)	+	+	+
Vitrella illyrica (A.J. Wagner, 1907)	+	+	+
Vitrella pygmaea (O. Boettger, 1880) <sup>127</sup>	—	+	—
Vitrella selecta L. Pintér, 1972	+	+	+
Vitrella sturanyi (A.J. Wagner, 1907)	+	+	* <sup>128</sup>
Vitrella subrimata (Reinhardt, 1871)	+	+	+
Gyralina circumlineata (L. Pfeiffer, 1846)	+	+	+
Gyralina gjirokastrana Riedel & Welter-Schultes, 1996	+	+	+
Gyralina candida candida (A.J. Wagner, 1909)	+	+	+
Gyralina nopscai Riedel, Feher & Eröss, 1999	+	+	—
Gyralina lunikense sp. n. <sup>129</sup>	—	—	—
Gyralina tarabosensis (Riedel, 1970)	+	+	+
Gyralina korabensis (Riedel, 1970)	—	+	* <sup>130</sup>
Aegopinella minor (Stabile, 1864)	+	+	+
Oxychilus camelinus (Bourguignat, 1852)	+	+	+

<sup>122</sup> Mentioned as *S. pantocratoris splendens*.

<sup>123</sup> Mentioned as *P. papillaris*.

<sup>124</sup> "does not live in today's Albania"

<sup>125</sup> "occurrence in Albania needs confirmation"

<sup>126</sup> First Albanian record: Reischütz et al. (2008b)

<sup>127</sup> First Albanian record: Fehér & Eröss (2009).

<sup>128</sup> "occurrence in Albania needs confirmation"

<sup>129</sup> Described in the Fehér & Eröss (2009).

<sup>130</sup> "Lives probably in today's Albania"

[Schistophallus camelinus]			
Oxychilus cyprius (L. Pfeiffer, 1847) [Schistophallus cyprius]	+	+	+
Oxychilus depressus (Sterki, 1880) [Mediterranea depressa]	+	+	+
Oxychilus hydatinus (Rossmässler, 1838) [Mediterranea h. hydatina]	+	+	+
Oxychilus juliae Riedel, 1990 [Mediterranea juliae]	+	+	+ <sup>131</sup>
Oxychilus planorbis (Möllendorff, 1899) [Mediterranea planorbis]	+	+	
Oxychilus inopinatus (Ulicný, 1887) <sup>132</sup> [Mediterranea inopinata]	—	+	—
Oxychilus glaber (Rossmässler, 1835)	[Morlina glabra nitidissima (Mousson, 1859)]	+	+
	[Morlina glabra striaria (Westerlund, 1881)]	+	+
Oxychilus draparnaudi (H. Beck, 1837)	+	+	+
Carpathica stussineri (A.J. Wagner, 1895)	+	+	+
Daudebardia brevipes (Draparnaud, 1805)	+	+	+
Daudebardia rufa (Draparnaud, 1805)	+	+	+
Paraegopis albanicus (Rossmässler, 1836)	+	+	+
Paraegopis bizonus A.J. Wagner, 1914	+	+	+
*Paraegopis mauritii (Westerlund, 1886)	—	+	* <sup>133</sup>
Paraegopis skipetaricus A.J. Wagner, 1914	+	+	+ <sup>134</sup>
*Aegopis laughofferi A.J. Wagner, 1914 <sup>135</sup> [A. septentrionalis laughofferi] <sup>136</sup>	+	+	* <sup>137</sup>
*Aegopis acies (Férussac, 1832) <sup>138</sup>	—	+	* <sup>139</sup>
*Aegopis verticillus (Lamarck, 1822) <sup>140</sup>	—	+	* <sup>141</sup>
Allaegopis amphikypellon Riedel, 1982	+	+	+
Allaegopis skanderbegianus (Polinski, 1924)	+	+	+
*Allaegopis transiens (Mousson, 1859) <sup>142</sup>	—	+	*
Vitrinidae			
Oligolimax annularis (S. Studer, 1820)	+	+	+
Oligolimax apatelus (Soós, 1924)	+	+	+ <sup>143</sup>
Vitrina pellucida (O.F. Müller, 1774)	+	+	+
Semilimacella bonellii reitteri (O. Boettger, 1880)	+	+	+
*Semilimacella carniolica (O. Boettger, 1884)	+	+	* <sup>144</sup>
Milacidae			
Tandonia albanica (Soós, 1924)	+	+	+
Tandonia sowerbyi (A. Férussac, 1823)	+	+	+
Limacidae			

<sup>131</sup> Listed together as "complex"

<sup>132</sup> First Albanian record: Fehér & Erőss (2009).

<sup>133</sup> "occurrence in Albania needs confirmation"

<sup>134</sup> "taxonomic revision is needed"

<sup>135</sup> Most probably does not live in Albania.

<sup>136</sup> According to Fauna Europaea, the nominate subspecies of this taxon is *A. septentrionalis* (Kobelt, 1899).

Kobelt's name, however, can not be used, because that is a junior homonym of *Zonites septentrionalis* Bourguignat, 1870. Thus probably *laughofferi* is the oldest available name for the species.

<sup>137</sup> "taxonomy and distribution not clear"

<sup>138</sup> Most probably does not live in Albania.

<sup>139</sup> "occurrence in Albania needs confirmation"

<sup>140</sup> Most probably does not live in Albania.

<sup>141</sup> "occurrence in Albania needs confirmation"

<sup>142</sup> "occurrence in Albania needs confirmation"

<sup>143</sup> "systematic position not clear"

<sup>144</sup> "systematic position not clear"

Lehmannia marginata (O.F. Müller, 1774)	+	+	+
Lehmannia szigethyae Wiktor, 1975	+	+	+
Lehmannia nyctelia (Bourguignat, 1861) <sup>145</sup>	—	+	—
Limacus flavus (Linnaeus, 1758)	+	+	+
Limax cephalonicus Simroth, 1886	+	+	+
Limax cinereoniger Wolf, 1803	+	+	+
Limax conemenosi O. Böttger, 1882 <sup>146</sup>	—	+	—
Limax graecus Simroth, 1889 <sup>147</sup>	—	+	—
Limax maximus Linnaeus, 1758	+	+	+
Limax wohlberedti Simroth, 1900	+	+	+
Malacomimax mrazekii (Simroth, 1904) <sup>148</sup>	—	+	—
Agriolimacidae			
Deroferas agreste (Linnaeus, 1758)	+	+	+
Deroferas reticulatum (O.F. Müller, 1774)	+	+	+
*Deroferas sturanyi (Simroth, 1894)	+	+	* <sup>149</sup>
Deroferas turicum (Simroth, 1894)	+	+	+
*Gigantomilax csikii Soós, 1924	—	—	+ <sup>150</sup>
Arionidae			
Arion subfuscus (Draparnaud, 1850)	+	+	+
[A. fuscus (O.F. Müller, 1774)]			
Arion circumscriptus Johnston, 1828	+	+	+
Arion fasciatus (Nilsson, 1823) <sup>151</sup>	—	+	—
*Arion rufus (Linnaeus, 1758)	—	+	* <sup>152</sup>
Helicodontidae			
Lindholmiola corycensis (Rossmässler, 1838)	+	+	+ <sup>153</sup>
Lindholmiola lens (Férussac, 1832)	—	+	+
*Helicodonta laughofferi A.J. Wagner, 1912	+	+	* <sup>154</sup>
Helicodonta obvoluta albanica A.J. Wagner, 1914	+	+	+
Cochlicellidae			
Cochlicella acuta (O.F. Müller, 1774)	+	+	+
Cochlicella barbara (Linnaeus, 1758)	+	+	+
Hygromiidae			
Xeropicta derbentina (Krynicki, 1836)	+	+	+
Semifruticicola serbica A.J. Wagner, 1914 [Semifruticicola costulata serbica A.J. Wagner, 1914] <sup>155</sup>	+	+	* <sup>156</sup>

<sup>145</sup> First Albanian record: Fehér & Erőss (2009).

<sup>146</sup> First Albanian record: Fehér & Erőss (2009).

<sup>147</sup> First Albanian record: Fehér & Erőss (2009).

<sup>148</sup> First Albanian record: Fehér & Erőss (2009).

<sup>149</sup> "occurrence in Albania needs confirmation"

<sup>150</sup> "probably belongs to *Deroferas*, but doubtful"

<sup>151</sup> First Albanian record: Fehér & Erőss (2009).

<sup>152</sup> "Lives probably not in today's Albania"

<sup>153</sup> Dhora & Welter-Schultes (1996) refer to this species as *Lindholmiola girva* (Frivaldszky, 1835), because they, as well as e.g. Irikov & Mollov (2006) treat these names as synonyms. In Subai's opinion – which is supported by molecular phylogenetic data (see Fehér & Erőss 2009) – *L. girva* and *L. corycensis* are morphologically distinguishable, distinct species; and only the latter one occurs in Albania.

<sup>154</sup> "Lives probably not in today's Albania"

<sup>155</sup> According to Fauna Europaea, the nominate subspecies of this taxon is *Helix costulata* Brancsik, 1897, however that is a homonym of *Helix costulata* Pfeiffer, 1828. Therefore *serbica* seems to be the oldest available name for the species.

<sup>156</sup> "systematic position not clear"

Trochulus erjaveci (Brusina, 1870) <sup>157</sup> [Trichia erjaveci]	+	+	+
Candidula castriota Soós, 1924	+	+	+ <sup>158</sup>
Candidula rhabdotoides (A.J. Wagner, 1928) <sup>159</sup>	—	+	—
Xerotricha conspurcata (Draparnaud, 1801)	+	+	+
Cernuella cisalpina (Rossmässler, 1837) <sup>160</sup>	+	+	—
Cernuella virgata (Da Costa, 1778) <sup>161</sup>	+	+	+
*Hiltrudia mathildae (Westerlund, 1881) <sup>162</sup>	+	+	+
Hiltrudia kusmici (Clessin, 1887) <sup>163</sup>	—	+	—
Metafruticicola occidentalis Subai, 1999	+	+	—
Monachoides fallax (A.J. Wagner, 1914)	+	+	—
Monachoides incarnatus (O.F. Müller, 1774)	+	+	+
Pseudotrichia rubiginosa (Rossmässler, 1838) <sup>164</sup>	+	+	+
Xeromunda vulgarissima (Mousson, 1859)	+	+	+
Xerolenta obvia (Menke, 1828)	+	+	+
Xerocrassa muehlfeldtiana (Rossmässler, 1837)	+	+	— <sup>165</sup>
Trochoidea pyramidata (Draparnaud, 1805)	+	+	+
Trochoidea trochoides (Poirier, 1789)	+	+	+
Monacha cartusiana (O.F. Müller, 1774)	+	+	+
Monacha subobstructa (Bourguignat, 1855) [Monacha claustralis (Menke, 1828)] <sup>166</sup>	+	+	+ <sup>167</sup>
Monacha cf. dofleini (P. Hesse, 1928) <sup>168</sup>	—	+	—
Monacha emigrata sattmanni Hausdorf, 1996	+	+	+
Monacha emigrata senitshika Hausdorf, 1996	+	+	+
Monacha frequens (Mousson, 1859)	+	+	+
Monacha microtricha S.H.F. Jaeckel, 1954	+	+	* <sup>169</sup>
Monacha parumcincta (Menke, 1828)	+	+	* <sup>170</sup>
Helicidae			
Theba pisana pisana (O.F. Müller, 1774)	+	+	+
Helix cincta O.F. Müller, 1774	+	+	* <sup>171</sup>
Helix dormitoris Kobelt, 1898	+	+	+

<sup>157</sup> Dhora & Welter-Schlüter (1996) indicate the subspecies *T. e. floerickei* (Kobelt, 1898) for Albania, which is treated as a synonym by Fauna Europaea.

<sup>158</sup> "systematic position is not clear"

<sup>159</sup> First Albanian record: Fehér & Erőss (2009).

<sup>160</sup> According to Fauna Europaea, this is the senior synonym of *C. jonica* (Mousson, 1854), and therefore listed for Albania. See also the note for *C. virgata*.

<sup>161</sup> Dhora & Welter-Schlüter (1996) treat as the senior synonym of *C. jonica* (Mousson, 1854).

<sup>162</sup> This species does not live in Albania. See notes for *H. kusmici*.

<sup>163</sup> See notes and some Albanian records in the Fehér & Erőss (2009).

<sup>164</sup> First verified Albanian record: Fehér & Erőss (2009).

<sup>165</sup> "indeterminate sp."

<sup>166</sup> According to F. Welter-Schlüter (personal communication) the name *claustralis* can not be used for this species. Although Hausdorf (2000: 82) argued that Mousson (1854, describing a species from Greece) "adopted" the name *claustralis* from Menke (1828, published in synonymy) and used it for a valid species (Code Art. 11.6), this was incorrect. Mousson (1859: 254) described a new species and had no idea that Menke (1828: 12) had mentioned such a name as a synonym in a list of names. This species should then be called *Monacha subobstructa* (Bourguignat, 1855), because this is the next available name according to Hausdorf (2000).

<sup>167</sup> Mentioned as *M. dissimilans* Pintér, 1968.

<sup>168</sup> First mentioned for Albania by Reischütz et al. (2008b)

<sup>169</sup> "systematic position not clear"

<sup>170</sup> "occurrence in Albania needs confirmation"

<sup>171</sup> "occurrence in Albania needs confirmation"

<i>Helix lucorum</i> Linnaeus, 1758	+	+	+
<i>Helix philibinensis</i> Rossmassler, 1839	—	+	* <sup>172</sup>
* <i>Helix oestreichi</i> Kobelt, 1902	+	+	
<i>Helix secernenda</i> Rossmässler, 1847	+	+	+
<i>Helix vladika</i> (Kobelt, 1898) [H. vladica]	+	+	+
<i>Cornu aspersum</i> (O.F. Müller, 1774)	+	+	+
<i>Eobania vermiculata</i> (O.F. Müller, 1774)	+	+	+
<i>Cepaea vindobonensis</i> (C. Pfeiffer, 1828)	+	+	+
<i>Codringtonia neocrassa</i> Zilch, 1952 <sup>173</sup>	+	+	* <sup>174</sup>
[ <i>Chilostoma apfelbecki bindzaensis</i> (Sturany, 1909)] <sup>175</sup>	+	+	+
<i>Cattania inflata</i> (Kobelt, 1876) [ <i>Chilostoma inflata</i> ]	+	+	+
<i>Cattania maranajensis</i> (A.J. Wagner, 1914) [ <i>Chilostoma maranajensis</i> ]	+	+	+
* <i>Cattania jakupicae</i> (Urbanski, 1979) [ <i>Chilostoma phocaea</i> <i>jakupicae</i> ] <sup>176</sup>	+	+	—
<i>Dinarica albanica</i> (S.H.F. Jaekel, 1954) [ <i>Chilostoma stenomphala albanica</i> ]	+	+	* <sup>177</sup>
<i>Dinarica petrovici</i> (A.J. Wagner, 1914) [ <i>Chilostoma petrovici</i> ]	+	+	+
<i>Dinarica pouzolzi montenegrina</i> (Rossmässler, 1836) [ <i>Chilostoma p. montenegrina</i> ]	+	+	+
* <i>Dinarica pouzolzi pouzolzi</i> (Deshayes, 1830) [ <i>Chilostoma p. pouzolzi</i> ] <sup>178</sup>	+	+	
<i>Dinarica serbica</i> (Kobelt, 1872) [ <i>Chilostoma serbica</i> ]	+	+	+
<i>Josephinella</i> (?) <i>korabensis</i> (Subai, 1997) <sup>179</sup>	—	—	—
<i>Josephinella</i> (?) <i>zebiana</i> (Sturany, 1907) [ <i>Chilostoma zebiana</i> ] <sup>180</sup>	+	+	+
<i>Josephinella byshekensis</i> (Knipper, 1941) [ <i>Chilostoma byshekensis</i> ]	+	+	+
*[ <i>Chilostoma edlaueri</i> (Knipper, 1941)] <sup>181</sup>	+	+	+
<i>Josephinella fuchsii</i> (Knipper, 1939) [ <i>Chilostoma fuchsii</i> ]	+	+	+
* <i>Josephinella pindica</i> (O. Boettger, 1886) [ <i>Chilostoma subzonata pindica</i> ]	+	+	+ <sup>182</sup>
<i>Liburnica albanograeca</i> (Subai, 1995) [ <i>Chilostoma albanograeca</i> ] <sup>183</sup>	—	+	* <sup>184</sup>
<i>Liburnica dochii</i> (Sturany, 1907) [ <i>Chilostoma dochii</i> ]	+	+	+
<i>Liburnica dunjana</i> (Knipper, 1941) [ <i>Chilostoma dunjana</i> ]	+	+	+
<i>Liburnica harpya</i> (Subai, 1995) [ <i>Chilostoma harpya</i> ]	+	+	+
<i>Liburnica hoffmanni walteri</i> (O. Boettger, 1886) [ <i>Chilostoma h. walteri</i> ]	+	+	+
<i>Liburnica setigera setigera</i> (Rossmässler, 1836) [ <i>Chilostoma s. setigera</i> ]	+	+	+

<sup>172</sup> "systematic position of *oestreichi* is not clear, possibly *H. secernenda* or *H. philibinensis*"

<sup>173</sup> First Albanian record: Dhora & Welter-Schultes (1999b).

<sup>174</sup> "lives possibly in South Albania"

<sup>175</sup> According to Subai (personal communication) its systematic position on the generic level is still doubtful.

<sup>176</sup> Most probably does not live in Albania.

<sup>177</sup> "systematic position not clear"

<sup>178</sup> According to Subai (personal communication) does not live in Albania.

<sup>179</sup> According to Subai (personal communication) the genus level position of this taxon is still doubtful.

<sup>180</sup> According to Subai (personal communication) the genus level position of this taxon is still doubtful.

<sup>181</sup> According to Subai (personal communication) this is a synonym for *J. byshekensis*.

<sup>182</sup> Due to a misinterpretation of Subai (1995), Dhora & Welter-Schultes (1996) erroneously included this species in the Albanian list. Most probably however, this species does not live in Albania.

<sup>183</sup> First Albanian record: Fehér & Erőss (2009).

<sup>184</sup> "probably lives in South Albania"

Superba grisea Subai & Fehér, 2006	—	—	—
Superba kulmakana Subai & Fehér, 2006	—	—	—
Superba skipetarica skipetarica (Subai, 1995) [Chilostoma skipetaricus]	+	+	+
Superba skipetarica asketa Subai & Fehér, 2006	—	—	—
*Superba reischuetzi Subai, 1990 [Chilostoma reschuetzi] <sup>185</sup>	+	+	+
*Arianta arbustorum (Linnaeus, 1758) <sup>186</sup>	+	+	—

## Literature

- ALBRECHT, C., TRAJANOVSKI, S., KUHN, K., STREIT, B. & WILKE, T. (2006): Rapid evolution of an ancient lake species flock: Freshwater limpets (Gastropoda: Aculyidae) in the Balkan lake Ohrid. — *Organisms Diversity & Evolution* 6: 294–307.
- ALBRECHT, C., SCHULTHEiß, R., KEVREKIDIS, T., STREIT, B. & WILKE, T. (2007): Invaders or endemics? Molecular phylogenetics, biogeography and systematics of *Dreissena* in the Balkans. — *Freshwater Biology* 52: 1525–1536. Singapore.
- BÁBA, K. (1982): Eine neue zoogeographische Gruppierung der ungarischen Landmollusken und die Wertung des Faunenbildes. — *Malacologia* 22 (1-2 [Proc. seventh intern. malac. Congress]): 441–454. Perpignan/Ann Arbor.
- BANK, R.A. & GITTEMBERGER, E. (1985): Notes on Azorean and European *Carychium* species (Gastropoda, Basommatophora, Ellobiidae). — *Basteria* 49: 85–100. Leiden.
- DHORA, DH. & WELTER-SCHULTES, F. W. (1996): List of species and atlas of the non-marine molluscs of Albania. — *Schriften zur Malakozoologie* 9: 90–197. Cismar/Ostholstein.
- DHORA, DH. & WELTER-SCHULTES, F. W. (1999a): Mollusc cenoses from different environments in Albania. — *Schriften zur Malakozoologie* 13: 61–66. Cismar/Ostholstein.
- DHORA, DH. & WELTER-SCHULTES, F. W. (1999b): Clausiliidae of Albania: recent finds and biogeographical evaluations (Gastropoda, Pulmonata). — *Biologia Gallo-hellenica* 25 (1): 15–24. Athens.
- ERÖSS, Z.P., FEHÉR, Z. & HUNYADI, A. (2005): Invasion of a North American Alien, *Planorbella anceps* (MENKE, 1830) (Mollusca: Gastropoda: Planorbidae), in the ancient Lake Prespa. — *Tentacle* 13: 6–7. Honolulu.
- ERÖSS, Z., FEHÉR, Z. & SZEKERES, M. (2006): New species taxa of *Montenegrina* O. BOETTGER, 1877 (Gastropoda: Pulmonata: Clausiliidae) from Albania and the neighbouring regions. — *Annalen der Naturhistorischen Museums in Wien* 107B: 181–208. Vienna.
- FALKNER, G., BANK, R. A. & PROSCHWITZ, T. VON. (2001): Check-list of the non-marine molluscan species-group taxa of states of northern, atlantic and central Europe (CLECOM I). — *Heldia* 4: 1–76.
- FEHÉR, Z., ERÖSS, Z. & VARGA, A. (2001): Contributions to the knowledge of the Albanian *Cochlostoma* fauna (Gastropoda: Cyclophoridae). — *Schriften zur Malakozoologie* 17: 71–78. Cismar/Ostholstein.
- FEHÉR, Z. (2004): A revision of the genus *Cochlostoma*, subgenus *Titanopoma* (Gastropoda, Caenogastropoda, Cochlostomatidae), in particular the forms occurring in Albania. — *Basteria* 68: 25–44. Leiden.
- FEHÉR, Z. & DRIMMER, L. (2004): Molluscs in the MEGYERI Collection. — *Soosiana* 31: 21–25. Nagykovácsi.
- FEHÉR, Z. & ERÖSS, Z. (2009) Contribution to the Mollusca fauna of Albania. Results of the field trips of the Hungarian Natural History Museum between 1992 and 2007. — *Schriften zur Malakozoologie* 25: 3–21. Cismar/Ostholstein.
- FEHÉR, Z., DELI, T. & SÓLYMOS, P. (submitted) Revision of *Granaria frumentum* (DRAPARNAUD, 1801) (Mollusca, Gastropoda, Chondrinidae) subspecies occurring in the eastern part of the species' range. — *Journal of Conchology*.
- GELEMBIUK, G. W., MAY, G. E. & LEE, C. E. (2006): Phylogeography and systematics of zebra mussels and related species. — *Molecular Ecology* 15: 1033–1050.
- HAUSDORF, B. (2000): The genus *Monacha* in Turkey (Gastropoda: Pulmonata: Hygromiidae). — *Archiv für Molluskenkunde* 128: 61–151. Frankfurt am Main.
- IRIKOV, A. & MOLLOV, I. (2006): Terrestrial gastropods (Mollusca: Gastropoda) of the Western Rhodopes (Bulgaria). In: BERON, P. (ed.) *Biodiversity of Bulgaria. 3. Biodiversity of Western Rhodopes (Bulgaria and Greece)*. I. Pensoft & Nat. Mus. Natur. Hist., Sofia, pp. 753–832.
- JAECKEL, S. H. F., KLEMM, W. & MEISE, W. (1957): Die Land- und Süßwasser-Mollusken der Nördlichen Balkanhalbinsel. — *Abhandlungen und Berichte aus dem Staatlichen Museum für Tierkunde in Dresden* 23: 141–205. Dresden.
- MEIER-BROOK, C. (1983): Taxonomic studies on *Gyraulus* (Gastropoda: Planorbidae). — *Malacologia* 24: 1–113. Philadelphia.

<sup>185</sup> Due to a misinterpretation of Subai (1995), Dhora & Welter-Schultes (1996) included this species in the Albanian list. Most probably however, this species does not live in Albania.

<sup>186</sup> Probably does not live in Albania.

- MENKE, C. T. (1828): Synopsis methodica molluscorum generum omnium et specierum earum, quae in museo Menkeano adservantur; cum synonymia critica et novarum specierum diagnosibus. pp. 91+ I-XII. Pyrmonti. (Gelpke).
- MOUSSON, A. (1859): Coquilles terrestres et fluviatiles, recueillies dans l'Orient par M. le DR. ALEX. SCHLÄFLI. — Vierteljahrsschrift der Naturforschenden Gesellschaft in Zürich 4: 12-36, 253-297. Zürich.
- NORDSIECK, H. (1970): Die *Chondrina*-Arten der dinarischen Länder. — Archiv für Molluskenkunde 100 (5-6): 243-261. Frankfurt am Main.
- NORDSIECK, H. (1971): Zur Anatomie und Systematik der Clausilien, IX. Dinarische Clausiliidae, III: Das Genus *Herilla*. — Archiv für Molluskenkunde 101 (1-4): 39-88. Frankfurt am Main.
- NORDSIECK, H. (1974): Zur Anatomie und Systematik der Clausilien, XV. Neue Clausiliien der Balkan-Halbinsel (mit taxonomischer Revision einiger Gruppen der Alopiinae und Baleinae). — Archiv für Molluskenkunde 104 (4-6): 123-170. Frankfurt am Main.
- NORDSIECK, H. (1995): Beiträge zur Nomenklatur der europäischen Binnenmollusken, VII. Kritische Anmerkungen und Berichtigungen von Arttaxa der Clausiliidae, 2. — *Hednia* 2 (3/4): 91-96. München.
- NORDSIECK, H. (2001): *Clausilia neutra* WESTERLUND 1898 (Gastropoda: Stylommatophora: Clausiliidae). — Mitteilungen der Deutsche Malakozoologische Gesellschaft, 66: 25-26. Frankfurt a Main.
- NORDSIECK, H. (2007): Worldwide doorsnails. Conchbooks, Hackenheim, 214 pp.
- NORDSIECK, H. (2008): *Alinda biplicata* (MONTAGU) and *Laciniaria plicata* (DRAPARNAUD) diversity in comparision, with the description of new subspecies. — Archiv für Molluskenkunde 137: 133-157. Frankfurt am Main.
- NORDSIECK, H. (2009) Revisory notes on the genus *Montenegrina* O. BOETTGER 1877 and description of new taxa: (Gastropoda: Stylommatophora: Clausiliidae). — Archiv für Molluskenkunde 138: 71-87. Frankfurt am Main.
- SOÓS, L. (1924): CSIKI ERNÖ állattani kutatásai Albániában. - Explorations zoologicae ab E. CSIKI in Albania peractae. XIII. Puhatestüek - Mollusca. In: A Magyar Tudományos Akadémia Balkán-kutatásainak tudományos eredményei. (P. TELEKI & E. CSIKI, eds. 1940) Vol. 1. CSIKI ERNÖ állattani kutatásai Albániában - Explorations Zoologicae ab E. CSIKI in Albania Peractae. Pars 2., pp. 177-194, The Hungarian Academy of Sciences, Budapest.
- REISCHÜTZ, A., REISCHÜTZ, P.L. & FISCHER, W. (2008a): Helleniká pantoía, 14: *Planorbarius corneus* "die Form des Grossen Prespa-Sees" (Makedonien, Nomos Florina, Griechenland) (Gastropoda: Pulmonata: Planorbidae). — Nachrichtenblatt der Ersten Voralberger Malakologischen Gesellschaft 15: 7-10. Rankweil.
- REISCHÜTZ, A., REISCHÜTZ, N. & REISCHÜTZ, P.L. (2008b) Ein Beirtag zur Kenntnis der Molluskenfauna Albaniens. — Nachrichtenblatt der Ersten Voralberger Malakologischen Gesellschaft 15: 37-39. Rankweil.
- REISCHÜTZ, A. & REISCHÜTZ, P.L. (2009) Es muß nich immer Griechenland sein: *Sciocochlea* C.R. BOETTGER 1935 (Gastropoda: Clausiliidae) aus Albanien, und andere malakologische Notizen. — Nachrichtenblatt der Ersten Voralberger Malakologischen Gesellschaft 16: 47-49. Rankweil.
- SUBAI, P. (1995): Beitrag zur Kenntnis albanischer und griechischer Ariantinae (Gastropoda: Helicidae). — Annalen der Naturhistorischen Museums in Wien 97B: 75-94. Vienna.
- SUBAI, P. (1999a): Beschreibung einer neuen Art *Argna thracica*, sowie Angaben über die griechischen Argnidae. — Schriften für Malakozoologie 13: 43-48. Cismar/Ostholstein.
- SUBAI, P. (2008): Revision of the Argninae, 1. The species of *Agardhiella* from the countries bordering the Adriatic Sea: (Gastropoda: Pulmonata: Pupillidae). — Archiv für Molluskenkunde 137: 75-103. Frankfurt am Main.
- SUBAI, P. (2009) New land snails from Montenegro and Albania: (Mollusca: Gastropoda). — Archiv für Molluskenkunde 138: 103-112. Frankfurt am Main.